

## STUDY ON BIOCHEMICAL CHANGES IN CHICKPEA (*CICER ARIETINUM L.*) GENOTYPES DURING HIGH-TEMPERATURE STRESS

MANISH KUMAR SINGH

Directorate of Education, ICFRE, Dehradun Email: kumar.manish21@yahoo.co.in

Received - 23.11.2017; Reviewed and accepted - 06.12.2017

### ABSTRACT

**Objective:** Plants experience high temperature in many different ways, and adaptation or acclimation to high temperature occurs over different levels of plant organization. Keep this in mind the present work was undertaken to study the effect of high-temperature stress on antioxidant enzymes in chickpea genotypes. **Material Used** Four chickpea genotypes differing in their sensitivity to high-temperature stress were taken and grown in growth chambers in the phytotron facility of IARI, New Delhi. The plants were maintained at 18/23°C (control) and 25/35°C (temperature stress) night/day temperature after maximum tillering. **Results:** In all chickpea genotypes high-temperature stress increased membrane injury index (MI), the activity of superoxide dismutase (SOD) and ascorbate peroxidase (APX), malic acid and slightly decreased the activity of glutathione reductase (GR), relative water content (RWC), chlorophyll and carotenoid contents. Under the HT stress, the tolerant genotypes Pusa-1103 and BGD-72 exhibited higher RWC, chlorophyll and carotenoid, the activity of SOD, APX, GR and, and less decrease in MI as compared to susceptible genotypes Pusa-256 and RSG-991. Antioxidant enzymes showed a positive correlation (*r*) with chlorophyll content, RWC and negative with MI under high-temperature stress. **Conclusion:** From the results, it is apparent that the antioxidant defense mechanism plays an important role in heat stress tolerance in chickpea genotypes and thus screening of tolerant genotypes by antioxidant enzymes, would be helpful in a further breeding programme of Chickpea.

**Keywords:** Antioxidant enzymes, high-temperature stress, membrane injury index, chlorophyll contents, relative water content.

### INTRODUCTION

Chickpea (*Cicer arietinum L.*) is the third most important pulse crop in the world after dry beans and dry pea in the global farming community and is widely cultivated in West and South Asia and North African countries. The latest estimate for 2007-08 indicates that the production of pulses in India is 16.64 million tonnes from an area of 26.15 million hectares. In spite of having largest area under chickpea in the world, India's position in average productivity is yet to see a breakthrough to meet the per capita availability of 50 g pulses/day to alleviate proteins energy malnutrition.

In northern India, however, late planting of chickpea is done after harvest of rice, early potato or cotton. Such late sown chickpea crop experiences high temperature at the end of the cropping season. This high temperature at the end of cropping season led to the problem of poor biomass and forced maturity [8]. The Inter-Governmental Panel on Climate Change (IPCC) of the United Nations in its recent report has confirmed the global warming trends and projected that the globally averaged temperature of the air above the earth's surface would rise by 1.4-5.8 °C over the next 100 years [18].

This high-temperature stress leads to the production of reactive oxygen species (ROS) which damages the plant cellular and subcellular system. However, plants protect its systems from toxic effects of the reactive oxygen species using antioxidant enzymes such as superoxide dismutase, ascorbate peroxidase, glutathione reductase, catalase, ascorbic acid and carotenoids [26]. Use of ion leakage and relative water content as simple indices for screening genotypes against heat and drought stress in chickpea and wheat has been suggested by many workers [9,24].

The present work was conducted to study the effect of high-temperature stress on antioxidant enzymes in chickpea genotypes.

### MATERIALS AND METHODS

The experiment was conducted at phytotron facility of I.A.R.I., New Delhi, with four chickpeas (*Cicer arietinum L.*) genotypes differing in sensitivity to high temperature (HT) stress, i.e., Pusa-1103 and BGD-72 (HT tolerant), Pusa-256 and RSG-991 (HT susceptible). Seeds of these four selected genotypes were collected from Genetics Division, I.A.R.I., New Delhi. They were treated with *Mesorhizobium ciceri* SPG strain and were sown in

earthen pots (20 × 30 cm<sup>2</sup>) containing a mixture of soil, sand and farmyard manures (FYM) in the ratio of 3:1:1. Recommended dose of nitrogen, phosphorus and potassium fertilizers were applied. Seeds were sown in each pot by dibbling method at 2 cm depth. Thinning was done at ten days after sowing, and five plants were retained in each pot.

Chickpea plants were exposed to temperature stress by covering them with polyvinyl chloride sheets (Capri Hans, sunflex 0.15 mm thickness and transmittance 85 %) mounted on wooden structures of size 3 x 2 x 2 m. The wooden polythene chambers were kept 10 cm above the ground for circulation of air and to control the humidity inside the chamber. Thermometer was placed inside the poly cover and the level of temperature was recorded regularly. The temperature inside the poly cover was 6.1 °C higher than the ambient temperature. Twenty pots from each of the four genotypes were shifted inside the polycovers at the 78 DAS to expose plants to temperature stress. Thirty seven days after temperature stress, all the pots were taken out and kept under the normal environment (ambient temperature), and the physiobiochemical observations were recorded.

Relative water content in leaves was estimated according to the method described by [6]. Membrane injury index was estimated from all genotypes in three random replicates as suggested by [9]. Chlorophyll content was determined according to [16] and carotenoid content was estimated according to [21].

Enzyme extract for superoxide dismutase, ascorbate peroxidase, glutathione reductase and catalase was prepared by first freezing the weighed amount of leaf samples (1 g) in liquid nitrogen to prevent proteolytic activity followed by grinding with 10 ml extraction buffer (0.1 M phosphate buffer, pH 7.5, containing 0.5 mM EDTA and 1 mM ascorbic acid). Brie was passed through 4 layers of cheese cloth and filtrate was centrifuged for 20 min at 15000 rpm at 4 °C and the supernatant was used as enzyme.

Superoxide dismutase activity was estimated by recording the enzyme induced decrease in absorbance of formazone made by nitro-blue tetrazolium with superoxide radicals [11].

Ascorbate peroxidase activity was estimated by observing the decrease in absorbance due to ascorbic acid at 290 nm and the Glutathione reductase activity was assayed by recording the

increase in absorbance in the presence of oxidized glutathione and DTNB (5, 5-dithiobis-2-nitrobenzoic acid [27].

Pearson-product-moment correlation coefficient (r) between various antioxidant enzymes and total chlorophyll content, relative water content and membrane injury index was computed according to [13].

## RESULTS AND DISCUSSION

### Relative water content

The analysis of data (Table 1) showed that with the increase in temperature stress there was progressive decrease in the RWC of flag leaves. Under ambient temperature, the RWC was higher in BGD-72 and Pusa-1103 and less in Pusa-256 and RSG-991. However, under high temperature (HT) treatment Pusa-1103 and BGD-72 showed significantly higher RWC. Average decline in RWC under high-temperature stress was 14.21 %, Pusa-1103 showed lowest per cent decline (6.0 %), while RSG-991 showed the highest decline (18.99%) in RWC. Significant differences were also obtained between treatment and genotypic interaction.

### Membrane injury index

The membrane injury Index (MI) decreased under heat stress in all the genotypes (Table 2.). However, under ambient temperature (AT) the MI was lowest in Pusa-1103 (31.61 %) closely followed by BGD-72, and highest in Pusa-256. Under HT stress, the Pusa-1103 showed lowest MI and RSG-991 exhibited highest MI (46.47 %). The average increase in MI under high-temperature stress was 19.33 %, Pusa-1103 showed the lowest increase (9.80 %), while RSG-991 showed a highest increase (31.74%) in MI. The interaction between treatment and genotypes was also significant.

### Photosynthetic pigments

Data on Chlorophyll 'a', chlorophyll 'b', total chlorophyll, total carotenoid, chlorophyll a/b ratio are reported in (Fig.1,2,3,4 and 5). Chlorophyll 'a', chlorophyll 'b', total chlorophyll, total carotenoid and chlorophyll/ carotenoid ratio in leaves of chickpea genotypes were higher under AT condition, and a significant decline was observed for all the above parameters under HT condition. Pusa-1103 followed by BGD-72 exhibited significantly higher chlorophyll 'a', chlorophyll 'b', total chlorophyll, and chlorophyll a/b ratio compared to Pusa-256 and RSG-991 under both the conditions. Total carotenoid content was significantly higher in Pusa-256 under AT condition along with Pusa-1103 and BGD-72. There was a general decline of 15.87 % in carotenoid content in all genotypes under HT condition. Under HT condition RSG-991 showed higher reduction, while Pusa-1103 maintained comparatively higher carotenoid content. Under AT condition the chlorophyll/carotenoid ratio was higher in BGD-72 and Pusa-1103 closely followed by RSG-991 and lowest in Pusa-256. Under HT treatment, there was a nonsignificant reduction in chlorophyll/ carotenoid ratio, and among genotypes, Pusa-1103 possessed highest chlorophyll/ carotenoid ratio.

### Antioxidant enzyme activities

Superoxide dismutase (SOD) activity showed significant increase under HT compared to AT condition (Table. 3). Among the genotypes, BGD-72 showed significantly higher SOD activity followed by Pusa-1103 and Pusa-256. RSG-991 exhibited the lowest activity under HT condition. SOD activity increased by two folds in BGD-72 and by 5 % in RSG-991. A significant interaction was observed between treatment and genotypes.

APX increased significantly under HT condition compared to AT (Table 4). Under both, the conditions Pusa-1103 possessed higher APX activity, while RSG-991 showed the lowest activity. BGD-72 and Pusa-256 exhibited moderate activity under both the conditions.

Data on glutathione reductase (GR) activity is reported in (Table 5). GR activity decreased significantly under HT condition. Under AT condition Pusa-1103 showed higher activity followed by BGD-72. Under HT condition also Pusa-1103 and BGD-72 maintained a higher GR activity, while RSG-991 exhibited a greater decline. A

significant interaction was observed between treatment and genotypes for GR activity.

Results on Pearson-product-moment correlation coefficient (r) between various antioxidant enzymes, and total chlorophyll content, relative water content and membrane injury index are reported in (Table 6). The results revealed that under high-temperature condition there exist a significant positive correlation between antioxidant enzymes and chlorophyll content and RWC, and significant negative correlation with MI. However under ambient temperature nonsignificant correlation was observed between antioxidant enzymes and other parameters. Among the three antioxidant enzymes, the glutathione reductase showed significantly higher correlation (r) with other three physiological traits under high temperature condition.

Plants experience high temperature in many different ways and adaptation or acclimation to high temperature occurs over different levels of plant organization [20]. In this study the BGD-72 and Pusa-1103 maintained higher RWC, chlorophyll, carotenoid contents and lower MI under high-temperature condition than Pusa-256 and RSG-991. This shows that the tolerant genotype Pusa-1103 had a greater water retention capacity under HT stress. Due to high temperature induced higher transpiration situation similar to water stress is created and RWC becomes important under heat stress. Heat stress injury involves water deficit and cell turgor loss [2]. Maintenance of favorable water status is essential for plant's tolerance to heat stress [14, 19].

Heat stress-induced decrease membrane stability has been reported in faba bean leaf discs [10, 16] and [26] also reported that the tolerant genotypes possess lower membrane injury index and high RWC, which enable them to maintain better metabolic activities.


Leaf photosynthetic pigment content (chlorophylls and carotenoids) and pigment ratios, such as Chl a/b is good indicators for stress detection and tolerance [5] Chlorophyll 'a', chlorophyll 'b', total chlorophyll and total carotenoid contents in leaves of chickpea genotypes decreased, while an increase was observed in Chl a/b ratio under HT condition compared to AT condition. [7] also showed increase in the chlorophyll a/b ratio in stressed Nagcarlang tomato plants, suggesting that these relationships could be used as an indicator of tolerance and physiological status of the plants under stress condition.

Carotenoid protects the photosynthetic systems against singlet oxygen and also plays a pivotal role in the thermal dispersion of excess excitation energy [1, 4, 22, 28]. Higher carotenoid content in tolerant genotypes Pusa-1103 and BGD-72 signifies their tolerance capacity.


Tolerance to high-temperature stress in crop plants has been reported to be associated with an increase in antioxidant enzymes activity (5, 15, 25, 26, 29) reported significant increase in SOD activity under temperature stress in wheat genotypes, and a greater increase in tolerant genotype C 306, while the susceptible genotypes showed lower activity. In this study the chickpea tolerant genotypes BGD-72 and Pusa-1103 exhibited the higher activity of SOD, APX and GR compared to susceptible genotypes. This shows that the tolerant genotypes combated the ROS by maintaining efficient antioxidant mechanism. [3] also reported similar results in late sown heat-tolerant wheat varieties. Higher activity of various antioxidative enzymes in temperature tolerant genotypes of various crop species has also been reported by various workers [3,12, 23, 26].

Study of the correlation coefficient between various antioxidant enzymes with chlorophyll content, relative water content and membrane injury index at high temperature revealed a positive correlation with chlorophyll content, relative water content and negative correlation with membrane injury index. [3] also reported a significant positive correlation between chlorophyll content and antioxidant enzymes and a negative correlation between membrane injury index and antioxidant at high-temperature stress condition.


From the preceding discussion it is clear that exposure of chickpea genotypes to high-temperature stress for a medium duration of thirty-seven days, i.e., 78 to 115 DAS increased the activity of superoxide dismutase, ascorbate peroxidase, and glutathione reductase. The temperature-tolerant genotypes Pusa-1103 and BGD-72 exhibited a comparatively higher superoxide dismutase, ascorbate peroxidase and slight decline in glutathione reductase compared to susceptible genotypes Pusa-256 and RSG-991. Efficient antioxidant enzymes status in tolerant genotypes under high-temperature condition reflected in lower membrane injury index, higher relative water content, chlorophyll and carotenoid content compared to susceptible genotypes Pusa-256 and RSG-991. Hence the selection of genotype based on these criteria may help in evolving chickpea genotypes tolerant to high-temperature stress with a better yield.




**Fig.1: Effect of high temperature stress on chlorophyll 'a' content in chickpea genotypes. Vertical bars show  $\pm$  S.E of mean. Data for treatments (T) and genotypes (G) and T x G interactions were significant (P=0.05)**




**Fig.2: Effect of high temperature stress on chlorophyll 'b' content in chickpea genotypes. Vertical bars show  $\pm$  S.E of mean. Data for treatments (T) and genotypes (G) and T x G interactions were significant (P=0.05)**



**Fig.3: Effect of high temperature stress on total chlorophyll content in chickpea genotypes. Vertical bars show  $\pm$  S.E of mean. Data for treatments (T) and genotypes (G) and T x G interactions were significant (P=0.05).**



**Fig.4: Effect of high temperature stress on total carotenoid content in chickpea genotypes. Vertical bars show  $\pm$  S.E of mean. Data for treatments (T) and genotypes (G) and T x G interactions were significant (P=0.05)**



**Fig.5: Effect of high temperature stress on chlorophyll 'a/b' content in chickpea genotypes. Vertical bars show  $\pm$  S.E of mean. Data for treatments (T).**

## REFERENCES

- Adams Demmig, B and Adams W W I (1992). Photo protection and other responses of plants to high light stress. *Ann. Rev. Plant Physiol Plant Mol. Biol.*, 43: 599–626.
- Ahmad S, Ahmad N, Ahmad R and Hamid M (1989). Effect of high-temperature stress on wheat productive growth. *J. Agric. Res. Lahore*, 27: 307–313.
- Almeselmani M, Deshmukh P S, Sairam R K, Kushwaha SR and Singh T P (2006). Protective role of antioxidant enzymes under high-temperature stress. *Plant Sci.*, 171: 382-388.
- Asada K, Endo T, Mano J and Miyake C (1998). Molecular mechanism for relaxation of and protection from light stress. Satan, K., Murata, N. (Eds). *Stress responses of photosynthetic organisms*. Amsterdam: Elsevier, pp. 37–52.
- Badiani M, Schenone G, Paolacci A R and Fumagalli I (1994). Daily fluctuations of antioxidant in bean (*Phaseolus vulgaris* L.) leaves. The influence of climatic factors, *Agrochimica*, 38: 25–36.
- Barrs H D and Weatherley (1962). A re-examination of the relative turgidity technique for estimating water deficit in leaves. *Aust. J. Biol. Sci.*, 15: 413-428.
- Camejo D, Rodri guez P, Morales M A, Dellamico J M, Torrecillas A, Alarcon J J (2005). High-temperature effects on the photosynthetic activity of two tomato cultivars with different heat susceptibility. *J. Plant Physiol.*, 162: 281–289.
- Chaturvedi S K and Dua R P (2003). Breeding chickpea for late sown conditions in northern India. *International Chickpea Conference*, Raipur, India, January, 20-22, pp. 11.
- Deshmukh P S, Sairam R K and Shukla D.S. (1991). Measurement of ion leakage as a screening technique for drought resistance in wheat genotypes. *Indian J. Plant Physiol.*, 34: 89-91.
- Deshmukh P S, Sairam R K, Kushwaha S R, Singh T P, Almeselmani M and Chaudhary, H.B. (2006). Physio-genetic approaches for increasing wheat productivity under

- rice-wheat cropping system. *Indian J. Agric. Sci.*, 76(11): 667-669.
11. Dhindsa R A, Plumb-Dhindsa P. and Thorpe T A (1981). Leaf senescence: Correlated with increased permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J. Exp. Bot.*, 126: 93-101.
  12. Gillham D J and Dodge A D (1987). Chloroplast superoxide and hydrogen peroxide scavenging systems from pea leaves: Seasonal variations. *Plant Sci.*, 50:105-109.
  13. Gomez K A and Gomez A A (1984). *Statistical Procedures for Agricultural Research*. 2nd Edn., John Wiley and Sons, Inc., New York, pp. 704.
  14. Graves W R, Joy R J and Dana M N (1991). Water use and growth of honey locust and tree-of-heaven at high root-zone temperature. *Hort. Sci.*, 26: 1309-1312.
  15. Gupta A S, Webb R P, Holaday A S and Allen R D (1993). Overexpression of superoxide dismutase protects plants from oxidative stress. Induction of ascorbate peroxidase in superoxide dismutase overexpression plants. *Plant Physiol.*, 103: 1067-1073.
  16. Hamada A M (2001). Alteration in growth and some metabolic processes of broad bean plants during extreme temperatures exposure. *Acta Physiol. Plant*, 23: 193-200.
  17. Hiscox J D and Israelstem G F (1979). A method for extraction of chlorophyll from leaf tissue without maceration using dimethyl sulfoxide. *Can. J. Bot.*, 57: 1332-1334.
  18. IPCC, 2001: *Climate change 2001: impacts, adaptation and vulnerability*, Contribution of Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change, edited by J. J. McCarthy, O. F. Canziani, N. A. Leary, D. J. Dokken and K. S. White (eds). Cambridge University Press, Cambridge, UK, and New York, USA, 2001.
  19. Jiang Y and Huang B (2000). Effects of drought or heat stress alone and in combination with Kentucky bluegrass. *Crop Sci.*, 40: 1358-1362.
  20. Larkindale J, Huang B (2005). Effects of abscisic acid, salicylic acid, ethylene and hydrogen peroxide in thermotolerance and recovery for creeping bentgrass. *Plant Growth Regul.*, 47: 17-28.
  21. Lichtenthaler H K and Wellburn A R (1983). Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents: *Biochem. Soc. Transactions*, 11: 591-592.
  22. Loggini B, Scartazza A, Brugnoli E and Navari-Izzo F (1999). Antioxidant defense system, pigment composition and photosynthetic efficiency in two wheat cultivars subjected to drought. *Plant Physiol.*, 119: 1091-1099.
  23. Menconi M, Sgherri C L M, Pinzino C, Navari-Izzo F (1995). Activated oxygen production and detoxification in wheat plants subjected to a water deficit programme. *J. Exp. Bot.*, 46: 1123-1130.
  24. Nagarajan S and Rane S (1997). Laboratory technique for quantifying heat tolerance. In: Nagarajan, S., Singh, H. and Tyagi, B.S. (Eds), *Wheat Research Needs Beyond 2000 AD*, Narosa Publishing House. New Delhi, India, pp. 209-226.
  25. Rui R L, Nie Y Q and Tong H Y (1990). SOD activity as a parameter for screening stress tolerant germplasm resources in sweet potato (*Ipomoea batatas* L.), *Jiangsu. J. Agric. Sci.*, 6: 52-56.
  26. Sairam R K, Srivastava G C and Saxena D C (2000). Increased antioxidant activity under elevated temperature: a mechanism of heat stress tolerance in wheat genotypes. *Biologia Plantarum*, 43: 245-251.
  27. Smith I K, Vierheller T L and Thorn C A (1988). Assay of glutathione reductase in crude tissue homogenates using 5, 5'-dithiobis (2-nitrobenzoic acid). *Anal. Biochem.* 175: 408-413.
  28. Young A (1991). The photo protective role of carotenoids in higher plants. *Physiol. Plant*, 83: 702-708.
  29. Zhau R G, Fan Z H, Li X Z, Wang Z W and Han W (1995). The effect of heat acclimation on membrane thermostability and relative enzyme activity. *Acta Agron.*, 21: 568-572